ELSEVIER

Environmental Pollution 132 (2004) I–IV

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 132

Number 1 2004

1 **Levels of PAHs in soil and vegetation samples from Tarragona County, Spain**
M. Nadal, M. Schuhmacher, J.L. Domingo

The levels of 16 polycyclic aromatic hydrocarbons (PAHs) were determined in soil and chard samples collected in an area with chemical and petrochemical industries.

13 **Determining the background levels of bismuth in tissues of wild game birds: a first step in addressing the environmental consequences of using bismuth shotshells**
R. Jayasinghe, L.J.S. Tsuji, W.A. Gough, J.D. Karagatzides, D. Perera, E. Nieboer

It is not clear whether or not bismuth is a nontoxic shot alternative.

21 **Phytoextraction of heavy metals by canola (*Brassica napus*) and radish (*Raphanus sativus*) grown on multicontaminated soil**
L. Marchiol, S. Assolari, P. Sacco, G. Zerbi

As anthropogenic soil pollution is usually not limited to single pollutants, plants potentially suited for phytoremediation should be observed in multicontaminated substrates.

29 **Accumulation of lead, zinc, copper and cadmium by 12 wetland plant species thriving in metal-contaminated sites in China**
H. Deng, Z.H. Ye, M.H. Wong

Metal accumulation by wetland plants differed among species, populations and tissues and is affected by substrate factors.

41 **Factors influencing liver PCB concentrations in sparrowhawks (*Accipiter nisus*), kestrels (*Falco tinnunculus*) and herons (*Ardea cinerea*) in Britain**
C.L. Wienburg, R.F. Shore

Over a six-year period in the 1990s, body condition, age and sex accounted for up to half of the intra-specific variation in liver PCB concentrations in predatory birds in Britain.

51 **Oribatid mite communities and metal bioaccumulation in oribatid species (Acari, Oribatida) along the heavy metal gradient in forest ecosystems**
P. Skubała, A. Kafel

Oribatid mites are positively stimulated by moderate concentration of heavy metals and they appear to be accumulators of copper.

61 **Microbial acidification and pH effects on trace element release from sewage sludge**
S. Qureshi, B.K. Richards, T.S. Steenhuis, M.B. McBride, P. Baveye, S. Dousset

The primary role of microbes in the potentially substantial mobilization of trace elements from wastewater sludge appears to be that of acidification.

73 **Fluxes of trichloroacetic acid through a conifer forest canopy**
R.T. Stidson, K.V. Heal, C.A. Dickey, J.N. Cape, M.R. Heal

Annualized within-canopy elimination of TCA is similar to that determined under controlled reductionist conditions.

85 **Growth of *Phragmites australis* (Cav.) Trin ex. Steudel in mine water treatment wetlands: effects of metal and nutrient uptake**
L.C. Batty, P.L. Younger

Growth of Phragmites australis is significantly reduced by metal toxicity in highly contaminated wetlands.

95 **Shade and flow effects on ammonia retention in macrophyte-rich streams: implications for water quality**
R.J. Wilcock, M.R. Scarsbrook, J.G. Cooke, K.J. Costley, J.W. Nagels

The interaction between macrophyte biomass and stream velocity determines ammonia removal capacity.

101 **Characterization of contaminants in snapping turtles (*Chelydra serpentina*) from Canadian Lake Erie Areas of Concern: St. Clair River, Detroit River, and Wheatley Harbour**
S.R. de Solla, K.J. Fernie

Organochlorine contaminant levels in some snapping turtle eggs in Canadian Lake Erie Areas of Concern exceed both minimum human consumption and Environmental Quality Guidelines.

S0269–7491(04)00308–2

113 Arsenic hyperaccumulation by *Pteris vittata* from arsenic contaminated soils and the effect of liming and phosphate fertilisation
N. Caille, S. Swanwick, F.J. Zhao, S.P. McGrath

Pteris vittata may be suitable for phytoremediation of soils moderately contaminated with arsenic.

121 Perchlorate in water, soil, vegetation, and rodents collected from the Las Vegas Wash, Nevada, USA
P.N. Smith, L. Yu, S.T. McMurry, T.A. Anderson

Concentrations of perchlorate in soil correlate with perchlorate concentrations in vegetation and rodents, but vegetation and rodent tissue concentrations are not highly correlated.

129 Cumulative impact of 40 years of industrial sulfur emissions on a forest soil in west-central Alberta (Canada)
J. Prietzel, B. Mayer, A.H. Legge

Forty years of industrial sulfur emissions caused accumulation of organic and inorganic sulfur in the soil, soil acidification, podzolization, and nutrient depletion in a boreal pine ecosystem.

145 White poplar (*Populus alba*) as a biomonitor of trace elements in contaminated riparian forests
P. Madejón, T. Marañón, J.M. Murillo, B. Robinson

White poplar leaves can be used as biomonitors of soil pollution by Cd and Zn.

157 Over one hundred years of trace metal fluxes in the sediments of the Pearl River Estuary, South China
C.C.M. Ip, X.D. Li, G. Zhang, J.G. Farmer, O.W.H. Wai, Y.S. Li

The combination of trace metal concentration, Pb isotopic composition, element normalisation and ^{210}Pb dating analysis can reveal the long-term accumulation of metals in estuarine sediments.

173 Spatial distribution of atmospheric PAHs and PCNs along a north–south Atlantic transect
F.M. Jaward, J.L. Barber, K. Booij, K.C. Jones

Proximity to source regions and air mass origin influence broad trends in oceanic air POPs concentrations, while diurnal cycling occurs in remote regions.

Number 2 2004

183 Accelerated remediation of pesticide-contaminated soil with zerovalent iron
P.J. Shea, T.A. Machacek, S.D. Comfort

Zerovalent iron promotes pesticide degradation in highly contaminated soil.

189 Sequential extraction of heavy metals in river sediments of an abandoned pyrite mining area: pollution detection and affinity series
F. Pagnanelli, E. Moscardini, V. Giuliano, L. Toro

A modified Tessier's procedure, discriminating organic and sulphide bound metals, was used to detect pollutant mobilisation from red mud to river sediment in an abandoned pyrite mine.

203 Selenium accumulation in the cockle *Anadara trapezia*
D.F. Jolley, W.A. Maher, J. Kyd

The marine bivalve Anadara trapezia is a good bioindicator for marine selenium contamination.

213 Irrigation runoff insecticide pollution of rivers in the Imperial Valley, California (USA)
V. de Vlaming, C. DiGiorgio, S. Fong, L.A. Deanovic, M. de la Paz Carpio-Obeso, J.L. Miller, M.J. Miller, N.J. Richard

Organophosphorous insecticides in runoff water from the USA and Mexico have impacted rivers in the Imperial Valley, California.

231 In situ and laboratory bioassays to evaluate the impact of effluent discharges on receiving aquatic ecosystems
R. Smolders, L. Bervoets, R. Blust

Responses of carp and zebra mussel to effluents differ under field and laboratory conditions.

245 An ecologically relevant exposure assessment for a polluted river using an integrated multivariate PLS approach
R. Smolders, W. De Coen, R. Blust

Multivariate analysis indicates how multiple lines of evidence can be integrated to provide a holistic overview of the impact of effluent discharges on the receiving aquatic ecosystem.

265 Modification of trace metal accumulation in the green mussel *Perna viridis* by exposure to Ag, Cu, and Zn
D. Shi, W.-X. Wang

Prior exposure of mussels to one metal may alter the rates of bioaccumulation of other metals.

279 Genotoxic and teratogenic potential of marine sediment extracts investigated with comet assay and zebrafish test
U. Kammann, S. Biselli, H. Hühnerfuss, N. Reineke, N. Theobald, M. Vobach, W. Wosniok

In vitro toxicity assays are used to assess genotoxic and teratogenic effects of environmental extracts.

289 Effects of heavy metals on growth and arsenic accumulation in the arsenic hyperaccumulator *Pteris vittata* L.
A.O. Fayiga, L.Q. Ma, X. Cao, B. Rathinasabapathi

Pteris vittata was effective in hyperaccumulating arsenic in the presence of heavy metals though at a reduced rate.

297 Growth response to ozone of annual species from Mediterranean pastures
B.S. Gimeno, V. Bermejo, J. Sanz, D. de la Torre, S. Elvira

The therophytes from dehesa acidic pastures of central of the Iberian peninsula present a great sensitivity to ozone, as derived from growth- and biomass-related variables.

307 The uptake of uranium by *Eleocharis dulcis* (Chinese water chestnut) in the Ranger Uranium Mine constructed wetland filter
R.A. Overall, D.L. Parry

Dissolved uranium in the surface water column is available for uptake by the aquatic macrophyte Eleocharis dulcis, which then translocates and stores it in its root system.

321 Effects of ozone on the foliar histology of the mastic plant (*Pistacia lentiscus* L.)
J. Reig-Armiñana, V. Calatayud, J. Cerveró, F.J. García-Breijo, A. Ibars, M.J. Sanz

Ozone causes alterations in the mesophyll, the conductive tissue and the secretory channels of the mastic plant (Pistacia).

333 Chelator induced phytoextraction and in situ soil washing of Cu
B. Kos, D. Leštan

Horizontal permeable barriers may be a more effective remediation technique for Cu.

341 Sediment quality in littoral regions of the Gulf of Cádiz: a triad approach to address the influence of mining activities
I. Riba, J.M. Forja, A. Gómez-Parra, T.Á. DelValls

An integrative assessment approach can be used to identify areas with relatively low levels of bioavailable pollutants in sediments.

355 A retrospective assessment of gold mining in the Reedy Creek sub-catchment, northeast Victoria, Australia: residual mercury contamination 100 years later
R.C. Churchill, C.E. Meathrel, P.J. Suter

Surface water Hg concentrations can exceed government guidelines.

365 Distinguishing sources of groundwater nitrate by ^{1}H NMR of dissolved organic matter
J. Lu, A.C. Chang, L. Wu

Dissolved organic matter has different signatures, depending on their sources.

Number 3 2004

375 Characterization of the organic matter of sludge: determination of lipids, hydrocarbons and PAHs from road retention/infiltration ponds in France
C. Durand, V. Ruban, A. Amblès, J. Oudot

Road runoff and urban sludge can be grossly contaminated with petroleum hydrocarbons, which may affect disposal options.

385 Immunotoxicology in wood mice along a heavy metal pollution gradient
K. Tersago, W. De Coen, J. Scheirs, K. Vermeulen, R. Blust, D. Van Bockstaele, R. Verhagen

Complex interactions among metal burden, immune response and parasite burden suggest negative effects of heavy metal exposure on the immune system of wood mice.

395 Soil pollution by a pyrite mine spill in Spain: evolution in time
J. Aguilar, C. Dorronsoro, E. Fernández, J. Fernández, I. García, F. Martín, M. Simón

Remediation measures carried out after the Aznalcóllar pyrite mine spill were effective in the reduction of the pollution, although three years after the accident many areas are still polluted by As.

403 Vegetative and structural characteristics of agricultural drainages in the Mississippi Delta landscapes
J.L. Bouldin, J.L. Farris, M.T. Moore, C.M. Cooper

Vegetated buffer areas provide effective mitigation for non-point source pollution from agriculture.

413 Fifty-year sedimentary record of heavy metal pollution (Cd, Zn, Cu, Pb) in the Lot River reservoirs (France)
S. Audry, J. Schäfer, G. Blanc, J.-M. Jouanneau

Reservoir sediments provide fine-scale information on the historical record of metal pollution in a watershed.

427 Role of exogenous and endogenous silicon in ameliorating behavioural responses to aluminium in a freshwater snail
A. Dobranskyte, R. Jugdaohsingh, E. Stuchlik, J.J. Powell, K.N. White, C.R. McCrohan

Toxicity of aluminium to freshwater snails is ameliorated by orthosilicic acid.

435 Effects of compost and phosphate on plant arsenic accumulation from soils near pressure-treated wood
X. Cao, L.Q. Ma

Compost amendment can reduce As exposure risk for humans by reducing As accumulation by vegetables and can be an important strategy for remediating CCA-contaminated soils.

443 Reproduction of mountain birch along a strong pollution gradient near Monchegorsk, Northwestern Russia
M.V. Kozlov, E.L. Zvereva

Variation in reproductive characteristics of mountain birch near the nickel–copper smelter at Monchegorsk is independent of pollution load.

453 Relations between metals (Zn, Pb, Cd and Cu) and glutathione-dependent detoxifying enzymes in spiders from a heavy metal pollution gradient
G. Wilczek, A. Babczyńska, M. Augustyniak, P. Migula

Glutathione-linked enzyme activity in spiders from polluted areas depends on hunting strategy and sex.

463 Characterization of aerosols and its radiative impacts over urban and rural environments—a case study from Hyderabad and Srisailam
K. Madhavi Latha, K.V.S. Badarinath

Atmospheric aerosol measurements at the urban and rural environments showed significant difference in radiative forcing.

469 Concentrations of ammonia and nitrogen dioxide at roadside verges, and their contribution to nitrogen deposition
J.N. Cape, Y.S. Tang, N. van Dijk, L. Love, M.A. Sutton, S.C.F. Palmer

Ammonia contributes as much as nitrogen dioxide to the deposition of nitrogen from vehicle exhausts to roadside vegetation.

479 Inhibition of plasma butyrylcholinesterase activity in the lizard *Gallotia galloti palmae* by pesticides: a field study
J.C. Sánchez-Hernández, R. Carbonell, A. Henríquez Pérez, M. Montealegre, L. Gómez

Chemical reactivation of lizard BChE activity is a suitable diagnostic method for evaluating field exposure to organophosphorus and carbamate pesticides.

489 Polychlorinated dibenzo-*p*-dioxins (PCDDs), dibenzofurans (PCDFs), biphenyls (PCBs), and polycyclic aromatic hydrocarbons (PAHs) and 2,3,7,8-TCDD equivalents (TEQs) in sediment from the Hyeongsan River, Korea
C.-H. Koh, J.S. Khim, K. Kannan, D.L. Villeneuve, K. Senthilkumar, J.P. Giesy

A combination of in vitro bioassay and chemical analysis identified that PCDDs/Fs are responsible for the dioxin-like activity present in the extracts of Hyeongsan River sediment.

503 Assessing plant response to ambient ozone: growth of young apple trees in open-top chambers and corresponding ambient air plots
W.J. Manning, D.R. Cooley, A.F. Tuttle, M.A. Frenkel, C.J. Bergweiler

Conditions in charcoal-filtered and non-filtered open-top chambers affected apple tree growth equally and prevented assessment of ambient ozone effects.

509 Ecological profiles of caddisfly larvae in Mediterranean streams: implications for bioassessment methods
N. Bonada, C. Zamora-Muñoz, M. Rieradevall, N. Prat

Ecological profiles of caddisfly larvae in Mediterranean streams using water quality variables.

523 The effect of soil composition and hydration on the bioavailability and toxicity of cadmium to hibernating juvenile American toads (*Bufo americanus*)
S.M. James, E.E. Little, R.D. Semlitsch

The effect of cadmium-contaminated soils on hibernating amphibians depends on cadmium concentration and soil composition and moisture.

533 Effects of heavy metal pollution on red wood ant (*Formica* s. str.) populations
T. Eeva, J. Sorvari, V. Koivunen

Five species of red wood ants vary in their sensitivity to heavy metal pollution but all of them had smaller colonies in a polluted area.

541 Determination of solid–liquid partition coefficients (K_d) for the herbicides isoproturon and trifluralin in five UK agricultural soils
C.M. Cooke, G. Shaw, C.D. Collins

Herbicide soil sorption described by a dual-phase adsorption model reflected soil partitioning, as influenced by soil OC and humic substances.